

Anusol[®] suppositories

(bismuth subgallate 59,0 mg, bismuth oxide 24,0 mg, zinc oxide 296,0 mg per suppository).
(Bismuth subgallate, bismuth oxide, zinc oxide).

Read all of this leaflet carefully because it contains important information for you.
ANUSOL SUPPOSITORIES are available without a doctor's prescription, for you to treat a mild illness.

- Nevertheless, you still need to use ANUSOL SUPPOSITORIES carefully to get the best results from it.
- Keep this leaflet. You may need to read it again.
- Do not share ANUSOL SUPPOSITORIES with any other person.
- Ask your health care provider or pharmacist if you need more information or advice.
- You must see your doctor if your symptoms worsen or do not improve after a few days.

What is in this leaflet

- What ANUSOL SUPPOSITORIES are and what it is used for
- What you need to know before you use ANUSOL SUPPOSITORIES
- How to use ANUSOL SUPPOSITORIES
- Possible side effects
- How to store ANUSOL SUPPOSITORIES
- Contents of the pack and other information

1. What ANUSOL SUPPOSITORIES are and what it is used for

- ANUSOL SUPPOSITORIES belong to a group of medicines in the form of suppositories and anal ointments.
- ANUSOL SUPPOSITORIES soothe the pain and irritation of piles and other related conditions around the anus and rectum.

2. What you need to know before you use ANUSOL SUPPOSITORIES

- Do not use ANUSOL SUPPOSITORIES:**
 - If you are allergic to any of the ingredients
 - If you have serious conditions of the rectum
 - If you are bleeding from the rectum, or if the condition carries on, consult your doctor.

Warnings and precautions

Take special care with ANUSOL SUPPOSITORIES:
Harmful effects to the nervous system may occur with prolonged or excessive use.

Children

It is not recommended to use ANUSOL SUPPOSITORIES in infants and children.

Other medicines and ANUSOL SUPPOSITORIES

Always tell your health care provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Pregnancy and breast-feeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before taking this medicine.

Driving and using machines

ANUSOL SUPPOSITORIES are not likely to affect your ability to drive or use any tools or machines.

3. How to use ANUSOL SUPPOSITORIES

Do not take by mouth.
The area around the anal opening should be cleansed with soap and warm water, then dried carefully by patting (not rubbing) with soft toilet tissue or cotton wool before the medicine is used.
The usual dose is as given below.

Adults:

Remove plastic cover and insert one suppository in the morning, one at bedtime and one after every bowel movement or as directed by your doctor.

- Tear along the perforation to remove one suppository from the strip (see figure 1 below)
- Grasp the two halves of the plastic cover firmly, holding between the thumb and forefinger of each hand (as indicated in figure 2 below). Gently pull apart so that the suppository is exposed.
- Take the suppository out of packaging and put it as deeply into the anus as possible.

The suppositories are specially shaped so that they slip easily into the anus. You may find it easier to put in the suppository if you place one foot on a chair or lie on your side with one leg drawn up as high as possible under the chin.



Figure 1

Figure 2

If you have the impression that the effect of ANUSOL SUPPOSITORIES is too strong or too weak, talk to your doctor or pharmacist.
Children: Should not be used in children.

⚠️ If you use more ANUSOL SUPPOSITORIES than you should
In the event of overdose, consult your doctor or pharmacist. If neither is available, seek help at the nearest hospital or poison control centre. Take along any medicine that is left and the carton so that the hospital can easily tell what medicine you have used.

⚠️ If you forget to use ANUSOL SUPPOSITORIES
Do not use a double dose to make up for forgotten individual doses.

4. Possible side effects

ANUSOL SUPPOSITORIES can have side effects.
Allergic reactions may occur. Stop using immediately if redness, irritation, swelling or pain continues or increases.
Not all side effects reported for ANUSOL SUPPOSITORIES are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking ANUSOL SUPPOSITORIES, please consult your health care provider for advice.
If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. You can also report side effects to SAHPRA via the "6.04 Adverse Drug Reaction Reporting Form", found online under SAHPRA's publications:
<https://www.sahpra.org.za/Publications/Index/8>.
By reporting side effects, you can help provide more information on the safety of ANUSOL SUPPOSITORIES.

5. How to store ANUSOL SUPPOSITORIES

- Store all medicines out of reach of children.
- Store at or below 30 °C, in a dry place
- Do not use after the expiry date stated on the label or carton. Return all unused medicine to your pharmacist. Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What ANUSOL SUPPOSITORIES contain
Each suppository contains the following active substances:
bismuth subgallate 59,0 mg, bismuth oxide 24,0 mg and zinc oxide 296,0 mg per suppository.
The other ingredients are balsam peru, fractionated coconut oil, hard fat, kaolin and titanium dioxide.

What ANUSOL SUPPOSITORIES look like and contents of the pack

White or off-white suppositories having the characteristic odour of Balsam Peru. Pack of 12 suppositories.

Holder of Certificate of Registration

MDI Healthcare CC
374 Anderson Street
Menlo Park, 0081
South Africa
Customer service: wecare@acdoco.co.za

This leaflet was last revised in

Date of last revision as approved by SAHPRA: 10 November 1997

Registration number

E/11.8/0915

Access to the corresponding Professional Information

An online copy of the PI is available at <http://acdoco.co.za/anusol/>

Name and address of manufacturer

Farma
10 Rue Bouché Thomas – ZAC d'Orgemont
BP 50723
49007, Angers Cedex
France

Registration details:

Botswana: S3; B9321890
Namibia: NS; 04/11.8/1522
Ghana: OT; FDB/SD.08-5196
Kenya: POM; 0416
Malawi: PIM; PMPB/PL 353/2
Nigeria: OT; 04-3089
Tanzania: POM; TAN 00,918 Do2A WAR
Uganda: POD; 1983/13/97
Zambia: P; 082/026
Zimbabwe: P; 98/16.3/3450

ANUS-4-101120PIL

Anusol[®] setpille

(bismutsubgallaat 59,0 mg, bismutoksied 24,0 mg, sinkoksied 296,0 mg per setpil).
(Bismutsubgallaat, bismutoksied, sinkoksied).

Lees die hele inligtingsblad noukeurig aangesien dit belangrike inligting vir u bevat.

- ANUSOL SETPILLE is beskikbaar sonder 'n dokter se voorskrif vir u om 'n matige siekte te behandel. Nietemin moet u ANUSOL SETPILLE versigtig gebruik om die beste resultate te kry.
- Hou hierdie inligtingsblad. U mag dit dalk weer moet lees.
- Moenie ANUSOL SETPILLE met enige ander persoon deel nie.
- Vra u gesondheidsorgverskaffer of apteker as u meer inligting of advies nodig het.
- U moet u dokter sien as u simptome vererger of nie verbeter na 'n paar dae nie.

Wat is in hierdie inligtingsblad

- Wat ANUSOL SETPILLE is en waarvoor dit gebruik word
- Wat u nodig het om te weet voor u ANUSOL SETPILLE gebruik
- Hoe om ANUSOL SETPILLE te gebruik
- Moontlike nuwe-effekte
- Hoe om ANUSOL SETPILLE te stoor
- Inhoud van die pak en ander inligting

1. Wat ANUSOL SETPILLE is en waarvoor dit gebruik word

- ANUSOL SETPILLE behoort aan 'n groep medisyne in die vorm van setpille en anale salwe.
- ANUSOL SETPILLE streef die pyn en irritasie van aambeie en ander verwante kondisies rondom die anus en rektum.

2. Wat u nodig het om te weet voor u ANUSOL SETPILLE gebruik

- Moet nie ANUSOL SETPILLE neem**
 - Indien u allergies vir enige van die bestanddele is nie
 - Indien u ernstige kondisies van die rektum het nie
 - Indien u van die rektum bloei nie, of indien die kondisie aanhou, konsulteer u dokter.

Waarskuwings en voorsorgmaatreëls

Neem spesiale sorg met ANUSOL SETPILLE:
Skadelike effektes op die senuweestelsel kan voorkom met langdurige of oormatige gebruik.

Kinders

Dit word nie aanbeveel om ANUSOL SETPILLE vir babas en kinders te gebruik nie.

Ander medisyne en ANUSOL SETPILLE

Vertel altyd jou gesondheidsorgverskaffer as jy enige ander medisyne neem. (Dit sluit alle komplementêre of tradisionele medisyne in.)

Swangerskap en borsvoeding

Indien u swanger is of borsvoed, dink u mag swanger wees of beplan om 'n baby te hê, raadpleeg asseblief u dokter, apteker, of ander gesondheidsorgverskaffer om advies voor u hierdie medisyne gebruik.

Bestuur en die gebruik van masjiene

Dit is onwaarskynlik dat ANUSOL SETPILLE jou vermoë om te bestuur of die gebruik van enige gereedschap of masjiene sal beïnvloed.

3. Hoe om ANUSOL SETPILLE te gebruik

Moet nie monddelings neem nie.
Die gebied om die anale opening behoort met seep en warm water gereinig te word en dan versigtig met sagte sneespapier of watte drooggetik te word (nie vryf nie) voordat medikasie gebruik word.
Die gewone dosis word hier onder aangedui.

Volwassenes:

- Verwyder plastiese omhulsel en plaas een setpil in soggens, een saans en een na elke ontlasting, of volgens voorskrif van u dokter.
- Skeur al langs die perforasie om een setpil uit die strokie te verwyder (sien skets 1 hieronder).
- Hou die twee helftes van die plastiese omhulsel styf tussen die duime en wysvingers van albei hande vas (soos aangedui in skets 2 hieronder). Trek versigtig uitmekaar totdat die setpil sigbaar is.
- Verwyder die setpil en steek dit so diep as moontlik in die anus op. Die setpille is spesifiek ontwerp sodat hulle maklik in die anus opgesteek kan word. U mag dit makliker vind om die setpille in te steek as u een voet op 'n stoel rus of op u sy lê en een been so hoog as moontlik onder u ken optrek.



Skets 1

Skets 2

As u onder die indruk is dat die effek van ANUSOL SETPILLE te sterk of te swak is, praat met u dokter of apteker.
Kinders: Behoort nie gebruik te word vir kinders nie.

⚠️ Indien u meer ANUSOL SETPILLE gebruik het as wat u moes
In geval van oordosering, raadpleeg u dokter of apteker. Indien een nie beskikbaar is nie, kry hulp by die naaste hospitaal of gifbeheersentrum. Neem enige medisyne wat oor is saam, asook die karton sodat die hospitaal maklik kan vasstel watter medisyne u geneem het.

⚠️ Indien u vergeet om ANUSOL SETPILLE te neem
Moenie 'n dubbeldosis neem om op te maak vir vergete individuele dosisse nie.

4. Moontlike nuwe-effekte

ANUSOL SETPILLE kan nuwe-effekte hê.
Allergiese reaksies kan voorkom. Hou dadelik op gebruik indien rootheid, irritasie, swelling of pyn voortduur of vererger.
Nie alle nuwe-effekte wat vir ANUSOL SETPILLE aangemeld is, is in hierdie voubljet ingesluit nie. Indien u algemene gesondheid agteruitgaan of u enige ongewenste reaksies ervaar terwyl u ANUSOL SETPILLE gebruik, raadpleeg asseblief u gesondheidsorgverskaffer vir advies.
Indien u enige nuwe-effekte waarneem wat nie in hierdie voubljet genoem word nie, stel asseblief u dokter of apteker in kennis.

Rapportering van nuwe-effekte

As jy nuwe-effekte kry, praat met jou dokter of apteker. Jy kan ook nuwe-effekte rapporteer aan SAHPRA via die "6.04 Adverse Drug Reaction Reporting Form", wat aanlyn gevind kan word onder SAHPRA se publikasies:
<https://www.sahpra.org.za/Publications/Index/8>.
Deur die rapportering van nuwe-effekte, kan jy help om meer inligting oor die veiligheid van ANUSOL SETPILLE te verskaf.

5. Hoe om ANUSOL SETPILLE te stoor

- Stoor alle medisyne buite bereik van kinders.
- Bewaar teen of benede 30 °C, in 'n droë plek.
- Moet nie na die vervaldatum wat op die etiket of karton aangedui word gebruik nie. Gee alle ongebruikte medisyne terug aan jou apteker. Moenie van ongebruikte medisyne ontslae raak deur dit in dreineringsstelsels of rioolsisteme (bv. toilette) te gooi nie.

6. Inhoud van die pak en ander inligting

Wat ANUSOL SETPILLE bevat
Elke setpil bevat die volgende aktiewe bestanddele: bismutsubgallaat 59,0 mg, bismutoksied 24,0 mg en sinkoksied 296,0 mg per setpil.
Die ander bestanddele is balsam peru, gefraksioneerde klapperolie, harde vet, kaolin en titaandoksied.

Hoe ANUSOL SETPILLE lyk en inhoud van die pak

Wit tot naaswit setpille met 'n kenmerkende balsamiese reuk.
Pakkie met 12 setpille.

Houer van die Registrasiesertifikaat

MDI Healthcare BK
Andersonstraat 374
Menlopark, 0081
Suid-Afrika
Kliëntediens: wecare@acdoco.co.za

Hierdie inligtingsblad was laas hersien in

Datum van laaste hersiening soos goedgekeur deur SAHPRA: 10 November 1997

Registrasienumer

E/11.8/0915

Toegang tot die ooreenstemmende professionele inligting

*n Aanlyn kopie van die PI is beskikbaar by <http://acdoco.co.za/anusol/>



GRAVITY // AMPLIFY // PHOTOTYPE

SMART ART PRODUCTION ART INFORMATION

PROJECT #: 2136154 Wave_1

SKU JOB #: 2136156

FILE NAME: PA2136156_R3.ai

CONTACT: Angela Pearson

E-MAIL: Angela.Pearson@olberdingbrands.com

Church & Dwight Co., Inc
500 Charles Ewing Boulevard
Ewing, NJ 08628
800.833.9532

MECHANICAL: ROUND 3

JOB INFORMATION

JOB DESCRIPTION: ANUSOL SUPPOSITORIES PIL ZA 2022

BRAND: Anusol

FRONT MATERIAL #: 508108 BACK MATERIAL #:

FRONT COPY CODE #: 25150 BACK COPY CODE #:

UPC #: 025150 LAST REVISION DATE: 02/15/2022

ADDITIONAL JOB INFORMATION (if available)

PRINT PROCESS: Litho PRINTER: MMP Packetis, CHAZELLES

SUBSTRATE: White paper PRINTER DIE#: 148x210mm

JOB SIZE: 25g (12s)

C&D APPROVALS

IP #:

FRONT COLORS

Cyan	Black				

BACK COLORS

Dieline

LIMITATION OF LIABILITY
DO NOT PLATE, ENGRAVE, OR PROCEED TO PRESS WITH THIS FILE WITHOUT PROPER INSPECTION
All files and proofs must be thoroughly inspected by the recipient immediately upon receipt. It is the responsibility of the recipient to examine and proofread all files and proofs. In the event file(s) and/or proofs are found to be incorrect upon inspection, call 513-281-0999 and ask for the separation contact. The liability of Olberding Brand Family shall be limited to correcting and replacing the materials that we supplied that were found to be defective. Olberding Brand Family will not be liable for any other loss or damage including, but not limited to, cost of Flexo printing plates that we did not make, print cylinders, inks, substrates/printed materials, and/or make ready and press time.

